



HALAL CONTROL

Cysteine Manufacturing from a Halal Perspective

**GIMDES 9th International Halal and Tayyib Conference ,
24.11.2017**

Dr. Ali Fanous

HALAL CONTROL




- Structure and properties of cysteine
- Biological function of cysteine
- Applications of cysteine
 - Food
 - Pharmaceutical products
 - Cosmetics
 - Animal feed
- Production methods of cysteine
 - Keratin Hydrolysate
 - Fermentation
 - Enzymatic Bioconversion

OUTLINE



Cysteine is a naturally occurring, sulfur-containing amino acid that is found in most proteins, although only in small quantities. Cysteine is unique amongst the amino acids as it contains a thiol group. When cysteine is oxidized it can form cystine, which is two cysteine residues joined by a disulfide bond.




STRUCTURE AND PROPERTIES



Due to the ability to undergo redox reactions, cysteine has antioxidant properties. Cysteine is required for biosynthesis of essential compounds including methionine, thiamine, Biotin and coenzyme A (1,2).

Cysteine also plays crucial roles in protein folding, assembly, and stability through disulfide-bond formation (3,2).




BIOLOGICAL FUNCTION OF CYSTEINE



In food industry, especially in bakery application, cysteine is used as flour additives to break up the gluten in flour, thus reducing its stickiness and facilitating the kneading of the dough.

One of the largest applications of cysteine is the production of various flavors. For example, reacting cysteine with sugars in a Maillard reaction yields meat flavors.



Cysteine reduces the disulfide bonds (S-S) within the gluten network (4).

APPLICATIONS OF CYSTEINE



In cosmetics area, cysteine is used in permanent hair wave preparation due to its ability to break disulfide bonds in keratin. The acetylated cysteine (N-acetylcysteine) is being used in the formulation of safe and effective products for antiaging and skin care products (5,6).

In pharmaceutical field, the cysteine derivative N-acetyl cysteine (NAC) is often used as a cough medicine.

Cysteine has been linked to aiding in the remedy of certain symptoms from drinking alcohol. It directly counteracts the poisonous effects of acetaldehyde, a particularly toxic by-product of alcohol in the human body.

APPLICATIONS OF CYSTEINE



As animal feed, cysteine is considered efficacious in partially meeting the requirements of Sulphur containing amino acids in animal (7).

Cysteine is required by sheep in order to produce wool. However, for sheep, it is an essential amino acid that cannot be synthesized by the sheep and must be taken in as food from grass. This means that during drought conditions, sheep produce less wool.



APPLICATIONS OF CYSTEINE



- Keratin Hydrolysate
- Fermentation
- Enzymatic Bioconversion

PRODUCTION METHODS OF CYSTEINE






Pig bristle




Human hair



Feathers

KERATIN HYDROLYSATE





* Critical points

KERATIN HYDROLYSATE



Advantage:

- Cheap and natural source, one ton of hair yields 100 kg of cysteine.

Disadvantage:

- The hair method requires 27 Kg of hydrochloric acid to produce just one kilogram of cysteine (8).
- Unpleasant odor.
- For pharmaceutical companies, it is crucial to eliminate the possibility of dangerous contaminations, such as the pathogens responsible for BSE, SARS or bird flu.
- Non-Halal raw materials can be used in the production.



KERATIN HYDROLYSATE



Advance technology is used to ferment the improved strain-producing cysteine that will produce high yields. New approach is used by optimizing the existing metabolism of the bacterium's native metabolism (*E.coli*).

This enables large quantity of cysteine to be released by the bacterium into the fermentation media, as 90 percent of pure cysteine ends up in the final product which fulfills the quality standards for foods and pharmaceutical industries (9).




FERMENTATION



E. coli cells synthesize L-cysteine using a simple two-step pathway.

- L-serine, the starting molecule for the pathway, is either acquired from the cell surroundings or made from other cell precursors.
- Acetyl-CoA is used to activate the L-serine to prepare it for incorporation of an atom of sulfur.



cysE serine acetyltransferase

cysK cysteine synthase A

cysM cysteine synthase B

Cysteine biosynthesis in *E. coli* (10)


FERMENTATION



Preparation of growth media



Strain collection



Carbohydrate

Protein (skim milk, peptone, yeast extract)


Trace elements (minerals, yeast extract)

Buffering (Minerals)

FERMENTATION



Cultivation of bacteria (fermentation)



FERMENTATION




Downstream processing (filtration, centrifugation, precipitation and crystallization).



FERMENTATION






FERMENTATION



L-serine is the precursor for L-cysteine; however it is very difficult to produce L-serine by direct fermentation. Thus, it requires metabolic engineering approach to produce L-cysteine by combining the overexpression of genes encoding for serinebiosynthetic enzymes and deletion of gene encoding for serine-degrading enzyme (11).



GM MICROORGANISM



No	Material	Source
1	Beef extract	Extracted from beef
2	Peptone	Hydrolysis of protein from meat, casein, or gelatin using acid or enzymes
	a. Acid-hydrolyzed casein	Hydrolysed protein from casein using acid for hydrolysis example: casamino acids (Difco); Hy-case (Sheffield)
	b. Enzymatic digest of casein	Hydrolysed casein using enzymes for hydrolysis Example: Casitone (Difco); Tryptone; trypticase peptone
	c. Enzymatic digest of meat protein	Hydrolysed protein from meat using enzymes for hydrolysis. Bactopeptone proteos pepton
	d. Enzymatic digest of plant (soy) protein	Hydrolysed protein from plant (soybean) using enzymes for hydrolysis. Example: Soyapeptone
3	Mixed hydrolysate	Biosate (yeast extract and casein digest); Trypton (mixed-enzymatic source)
4	Yeast extract	Extracted from yeast
5	Agar	Carbohydrate from sea weed

GROWTH MEDIA



- Proteins or hydrolyzed protein from animal (such as gelatin, meat) → CP : source of protein and enzymes for hydrolysis ?
- Hydrolisates of starchy material → CP : source of amylases enzymes for hydrolysis ?



ENZYMES AS PROCESSING AIDS IN MEDIA PREPARATION



Antifoam

Normally people use silicon, sulphonates, esters, and animal oil (lard) or fatty acids from animal oil.

Activated carbon

Activated carbon is usually obtained from wood, coconut shell, or animal bones.



Some bacteria, particularly in the genus of *Pseudomonas*, can hydrolyze DL-2-amino- Δ^2 -thiazoline-4-carboxylic acid (DL-ATC) to L-cysteine (12,13).


DL-ATC is a desirable substrate for product L-cysteine due to simplicity of the process for chemical synthesis of DL-ATC.

This method has the potential to produce high yield of cysteine with low energy requirement during production.



Enzymatic Bioconversion






DL-2-amino-4-thiazoline-4-carboxylic acid (DL-ATC)



Bioconversion by *Pseudomonas sp.*




Cysteine

Enzymatic Bioconversion



The conversion of DL-ATC to cysteine can be described in the following successive steps: (i) enzymatic reclamation of D-ATC to L-ATC; (ii) a ring-opening reaction of L-ATC to N-carbamyl-L-cysteine (L-NCC) as intermediate; and (iii) hydrolysis of L-NCC to L-cysteine (2, 14).



Selection medium

DL-ATC, **glycerol**, KH_2PO_4 , $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, $\text{MnSO}_4 \cdot 4\text{H}_2\text{O}$, agar and water (15).

Fermentation medium

DL-ATC, **yeast extract**, **peptone**, **meat extract**, NaCl, **glycerol** and distilled water (16).

Fermentation medium

DL-ATC, **glycerol**, **beef extract**, **yeast extract**, **peptone**, NaCl, $\text{MnSO}_4 \cdot \text{H}_2\text{O}$, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$. (16).

The pharmaceutical, cosmetic and food industries worldwide require up to 4,750 tons of cysteine per year (17).

Until recently, it was one of the few amino acids that had to be obtained from animal or human raw materials, for example human hair, feathers, pig bristles and hooves. With improvements in biotech, it has been possible to engineer a bacteria that produces excess cysteine.

Owing to the fact that many Haram materials can be used in the manufacturing process, cysteine is considered as a Halal critical material.

CONCLUSION



1. Wirtz M, Droux M, Hell R. O-acetylserine (thiol) lyase: an enigmatic enzyme of plant cysteine biosynthesis revisited in *Arabidopsis thaliana*. *J Exp Bot* 2004; 55: 1785-98.
2. Wada M, Takagi H. Metabolic pathways and biotechnology of L-cysteine. *Appl Microbiol Biotechnol* 2006; 73: 48-54.
3. Wirtz M, Droux M. Synthesis of the sulfur amino acids: cysteine and methionine. *Photosynth Res* 2005; 86: 345-62
4. https://www.wacker.com/cms/media/publications/downloads/6882_EN.pdf
5. Renneberg R. High grade cysteine no longer has to be extracted from hair. In Demain, AL, eds. *Biotechnology for beginners*. Academic Press: Amsterdam 2008, pp.106.
6. Hillebrand G, Bush RD. Use of N-acetyl-L-cysteine and derivatives for regulating skin wrinkles and/or skin atrophy. Great Britain Patent EP 0 734 718 A2. 1992.
7. European Food Safety Authority (EFSA). Scientific Opinion on the safety and efficacy of L-cystine for all animal species. EFSA Panel on Additives and Products or Substances used in Animal Feed (FEEDAP), Parma, Italy *EFSA Journal* 2013; 11: 3173
8. Nur Izzah Ismail NI, Hashim YZ , Jamal P, Rashidi Othman, Hamzah Mohd Salleh. Production of Cysteine: Approaches, Challenges and Potential Solution. *International Journal of Biotechnology for Wellness Industries*, 2014; 3: 95-101
9. Wacker Fermentation-Grade. Discovery a new dimension of pureness: vegetarian cysteine. Germany: Wacker Chemie AG. 2010
10. Matteo P. Ferla, Wayne M. Bacterial methionine biosynthesis. *Patrick- Microbiology* 2014; 160: 1571–158

REFERENCES



11. Peter-Wendisch P, Stolz M, Etterich H, Kennerknecht N, Sahn H, Eggeling L. Metabolic engineering of *Corynebacterium glutamicum* for L-serine production. *Appl Environ Microbiol* 2005; 71: 7139-7144.
12. Awano N, Wada M, Kohdoh A, Oikawa T, Takagi H, Nakamori S. Effect of cysteine desulfhydrase gene disruption on l-cysteine overproduction in *Escherichia coli*. *Appl. Microbiol. Biotechnol.* 2003, 62: 239-243.
13. Awano N, Wada M, Mori H, S Nakamori, Takagi H. Identification and Functional Analysis of *Escherichia coli* Cysteine Desulfhydrases. *Appl. Environ. Microbiol.*, 2005, 71: 4149-4152.
14. Shiba T, Takeda K, Yajima M, Tadano M. Genes from *Pseudomonas sp.* Strain BS involved in the conversion of L-2-amino-2-thiazolin-4-carbonic acid to L-cysteine. *Appl. Environ. Microbiol.* 2002, 68: 2179-2187.
15. Sano K, Yokozeki K, Tamura F, Yasuda N, Noda I, Mitsugi K. Microbial Conversion of DL-2-Amino-A2-Thiazoline-4-Carboxylic Acid to L-Cysteine and L-Cystine: Screening of Microorganisms and Identification of Products. *Applied and environmental microbiology.* 1977, 34: 806-810
16. Chen W, Chen X, Wang P. Enzymatic synthesis enhancement of mutagenised strain *Pseudomonas sp.*C-25 for L-cysteine production. *Journal of Chemical and Pharmaceutical Research.* 2014, 6:259-265
17. <https://www.wacker.com/cms/media/en/documents/feature-pdf/Cystein.pdf>

REFERENCES



Thank you for your attention

www.halalcontrol.de



HALAL CONTROL



www.halalcontrol.de